

# Perinatal Stress Effects on Later Anxiety and Hormone Secretion in Male Mandarin Voles

Feng Qin He, Gang Fang, Bo Wang, Xin Jun Guo, and Chun Lin Guo  
Xi'an University at Xi'an

The estradiol ( $E_2$ ), estrogen receptor- $\alpha$  ( $ER\alpha$ ), testosterone (T), and androgen receptor (AR) can contribute to anxiety, but whether they are associated with the reversion of prenatal adverse outcomes remains unclear. Here, we tested the interactive effects of prenatal maternal restraint stress and early postnatal short-term maternal separation on adult male mandarin vole (*Microtus mandarinus*) behavior and changes in  $E_2$ , T, and their receptors. The results showed that PS adult males (PS/NH) exhibited an increase in anxiety-like behavior in open-field and elevated plus-maze tests than the other 3 groups, including adult male offspring controls (PC/NH), adult male offspring controls with short-term maternal separation (PC/H), and PS adult males with short-term maternal separation (PS/H). The increase in anxiety-like behavior was associated with significantly lower  $E_2$  and T serum levels, had significantly more  $ER\alpha$  immunoreactive neurons ( $ER\alpha$ -IRs) in some brain regions, as well as significantly fewer AR immunoreactive neurons (AR-IRs) in some brain regions than the other 3 groups. We found it interesting that the PC/H and PS/H were similar to the PC/NH in that they did not produce anxiety-like behavior. However, early postnatal short-term maternal separation reversed prenatally induced changes in  $E_2$  and T serum levels and altered  $ER\alpha$ -IRs and AR-IRs in the brain. These data suggest that changes in anxious adults may be governed by early environmental factors and their interactions because changes in  $E_2$  and T serum levels and the distribution of  $ER\alpha$  and AR in the brain result in behavioral changes related to less anxiety into adulthood.

**Keywords:** androgen receptor, estrogen receptor- $\alpha$  ( $ER\alpha$ ), prenatal stress, estrogen ( $E_2$ ), testosterone (T)

Anxiety and depression are highly prevalent chronic and potentially life-threatening neurological disorders (Nestler et al., 2002). The hypothalamic–pituitary–adrenal (HPA) axis is a highly plastic and sensitive system, and dysregulation of the HPA axis causes anxiety (Van den Bergh, Van Calster, Smits, Van Huffel, & Lagae, 2008). The steroid hormones estradiol ( $E_2$ ) and testosterone (T) affect the HPA axis, and  $E_2$  and T respectively increase and attenuate adrenocorticotrophic hormone (ACTH) and corticosterone secretion (Franklin, Saab, & Mansuy, 2012). Therefore, abnormal secretion of estrogen and androgens can also contribute to mood disorders (Morrison, Brinton, Schmidt, & Gore, 2006). For example, several lines of evidence suggest that anxiety disorders are most frequent during times when estrogen levels are low and may be relieved by estrogen treatment (Best, Rees, Barlow, & Cowen, 1992; Sichel, Cohen, Robertson, Ruttenberg, & Rosenbaum, 1995; Arpels, 1996; Gregoire, Kumar, Everitt, & Studd, 1996). Lower

levels of testosterone and testosterone levels that decreased more slowly over the course of a day were related to higher levels of anxiety/depression symptoms and attention problems in adolescent boys (Granger et al., 2003). Low testosterone levels in hypogonadal and aging men have been associated with diagnosed depression (Eskelinen, Vahlberg, Isoaho, Kivelä, & Irjala, 2007). It has also been shown that T treatment in mice can decrease symptoms of anxiety (Zuloaga, Morris, Jordan, & Breedlove, 2008).

Prenatal stress (PS) as a result of maternal stress can result in an increased risk of anxiety and depression in offspring (Howerton & Bale, 2012), which can lead to changes in  $E_2$  and T levels and influence the HPA axis and regulate the activity and behavior of male offspring (Kapoor & Matthews, 2011; Tchernitchin, Tchernitchin, Mena, Unda, & Soto, 1999). The effect of estrogen on anxiety may be explained by estrogen causing changes in the estrogen receptor- $\alpha$  ( $ER\alpha$ ) in particular brain regions (Newhouse et al., 2010; Fedotova, 2013). The expression of  $ER\alpha$  is found in abundance in the medial amygdala, the bed nucleus of stria terminalis (BNST), and the preoptic area (Krezel, Dupont, Krust, Chambon, & Chapman, 2001).  $ER\alpha$  activation has been shown to increase fear and anxiety in male mice and rats (Morgan & Pfaff, 2001; Patisaul & Bateman, 2008). However, the relationship between the distribution of  $ER\alpha$  and anxiety disorders in other animals remains unreported except in mice and rats.

Male rat offspring born to mothers exposed to restraint stress during the last week of pregnancy have decreased plasma T levels, elevated basal ACTH levels, and blunted responses to stress (Richardson, Zorrilla, Mandyam, & Rivier, 2006). Thus, evidence suggesting a role for T in mediating the effects of PS on HPA-axis

This article was published Online First October 26, 2015.

Feng Qin He, Gang Fang, Bo Wang, Xin Jun Guo, and Chun Lin Guo, Genetic Engineering Laboratory, College of Biological and Environmental Engineering, Xi'an University at Xi'an.

This research was supported by the National Natural Science Foundation of China (Grant 61173113), the Science and Technology Research and Development Foundation of Shaanxi (Grant 2015JM3116), and Science and Technology Program of Xi'an (CXY1443WL09).

Correspondence concerning this article should be addressed to Feng Qin He, Genetic Engineering Laboratory, College of Biological and Environmental Engineering, Xi'an University at Xi'an, 1 Technology Six Road, Xi'an, 710065, China. E-mail: hefengqin68220@163.com

activity and behavior in male offspring is accumulating. Androgens play an important role in rodents by acting on androgen receptors (ARs) in key brain areas, such as the medial preoptic area (mPOA) and the medial amygdaloid nucleus (MeA; Bingham, Williamson, & Viau, 2006; Williamson & Viau, 2007). Under certain stressors, androgens may affect anxiety-related behavior by acting through ARs in brain regions such as the BNST, MeA, and hypothalamic paraventricular nucleus (PVN; Edinger & Frye, 2006).

The postnatal environment can also have a considerable impact on the development of an organism. To assess the effects of the postnatal environment on behavior and physiology, numerous studies have explored the effects of early separation from the mother on development of the offspring. Separation periods can range from a single separation during the preweaning period (Ellenbroek, van den Kroonenberg, & Cools, 1998) to repeated daily separations (15 min to 5 hr; Brake, Zhang, Diorio, Meaney, & Gratton, 2004; Meaney et al., 1991). In addition, separations can encompass almost the entire preweaning period (such as in artificial rearing; see Gonzalez, Lovic, Ward, Wainwright, & Fleming, 2001; Lovic & Fleming, 2004). On the basis of these temporal parameters, maternal separation studies have found multiple and opposing effects on behavior and physiology. Elevated indices of anxiety have been exhibited as a result of long periods of daily separation (Ellenbroek et al., 1998; Huot et al., 2002). In contrast, short periods of separation, or handling regimens have exhibited reduced separation anxiety responses in pups (Lehmann, Stöhr, & Feldon, 2000).

The mandarin vole (*Microtus mandarinus*) is a monogamous rodent that engages in pair bonding, affiliation, and biparental care (Guo et al., 2011; Tai, Wang, & Zhou, 2001). Social-bonding and pair-bonding formation are critical components of mental health (Young, Gobrogge, Liu, & Wang, 2011), and mandarin voles have emerged as a model for the study of social-emotional behaviors such as those associated with anxiety (He, 2014) and depression (He, Tai, Zhang, & Zhang, 2012). Previous experiments by our group have shown that PS (e.g., exposure to ethanol) increases anxiety and decreases central AR expression in adult mandarin voles (He, 2014).  $E_2$ ,  $ER\alpha$ , T, and AR can contribute to anxiety, but whether they are associated with the reversion of prenatal adverse outcomes is unclear. Therefore, the aims of the present study were to (a) determine whether a correlation exists between prenatal maternal restraint stress and the distribution of  $ER\alpha$  and ARs in brain regions associated with anxiety-like behavior; and (b) investigate whether early postnatal maternal separation can reverse these behaviors, and whether  $E_2$  and T serum levels and the distribution of  $ER\alpha$ -IRs and AR-IRs in the brain change accordingly.

## Method

### Animals

All experimental animals included were female and male mandarin voles whom we obtained from an outbred colony and fed at the College of Biotechnology, Xi'an University, Xi'an, PRC. Mandarin voles were captured from Lingbao voles in Henan, China, 1997. Adult females ( $N = 40$ , 30–36 g, 90 days old) were individually housed in clear plastic cages (30 × 20 × 15 cm) and

were maintained in a 14:10-hr light/dark cycle at 24–26 °C. Hardwood shavings and cotton were provided as substrate and bedding. Rabbit chow (Laboratory Animals Center, Xi'an Medical University, Xi'an, China), carrot, and malt were provided ad libitum. All methods were approved by the Institutional Animal Care and Use Committee, Xi'an University of Arts and Science. To avoid exogenous and endogenous environmental factors that might have induced differences between animals (e.g., infection, age, sisterhood), we captured new mandarin voles from Lingbao each year and chose healthy animals of similar mass. These new voles are paired with laboratory voles, and their offspring supplement our laboratory population to avoid inbreeding.

Monogamous female voles do not display spontaneous ovarian activity or ovulation in nonestrus (Sawrey & Dewsbury, 1985) and females are not receptive to males except when they are in estrus. However, estrus can be induced in females through estrogen administration alone (Carter, Witt, Auksi, & Casten, 1987). Once fertilization occurs, continued male presence significantly increases the probability that the pregnancy will be carried to term. Therefore, female mandarin voles were brought into estrus with estradiol benzoate (0.75 µg/g, 24 hr before testing) and progesterone (0.015 mg/g, 4–6 hr before testing) and we monitored the estrus state of females using vaginal smears. Each female in estrus was paired with an adult male whose bilateral or unilateral testes had descended ( $N = 40$ ; total 40 pairs). Pregnancy began when two ejaculations were observed (Day 0 of pregnancy; Ward, Ward, Denning, French, & Hendricks, 2002).

### Prenatal Stress Treatment

The gestation period in mandarin voles is about 21 days, as it is in rats and mice. The fetal and neonatal development of the testis of a sexually undifferentiated gonad involves a succession of events allowing the development of each testicular cell type (Olaso & Habert, 2000). In the rat, Sertoli cells differentiate and surround the germ cells to form the seminiferous cords from 13.5 to 14.5 days postconception (dpc), and then they proliferate until 3 weeks after birth (Pelliniemi, Frojman, & Paranko, 1993). Therefore, we speculate that testicular development occurs in the fetal male vole from 13.5 to 14.5 days postconception. In the present experiment, the period of maternal stress began after the development of the testes. Forty pregnant female mandarin voles were randomly assigned either to a PS or a PC group. Pregnant female voles in the PS group ( $N = 20$ ) were subjected to restraint stress during the last 2 weeks of pregnancy. The restraint-stress treatment involved removing the female from its home cage and placing it in a closed, narrow, transparent plastic tube (11 cm in length × 6 cm in diameter) for 30 min, 3 times a week, at different times in the morning. The control group ( $N = 20$ ) was left undisturbed in their cages. One to five pups were produced in each litter, and the sex ratio was 1.87 females to 1 male (He, Tai, et al., 2012; Tai, Wang, & Zhao, 1999). To exclude litter effects, only one male pup was chosen from each litter: The 20 male pups from the treatment group were denoted as PS, and the 20 male pups from the control group as prenatal control (PC). The extra pups in each litter were used in other experiments conducted by our group. Thus, these offspring were all from the dam's first litter in the present experiment, and dams were housed with sires throughout the prenatal and preweaning periods.

### Early Postnatal Short-Term Maternal Separation

The PS group was divided into two groups: PS with postnatal short-term maternal separation (i.e., handling; PS/H,  $N = 10$ ) and PS without postnatal short-term maternal separation (i.e., no handling; PS/NH,  $N = 10$ ). The PC group was divided into two groups: PC with postnatal short-term maternal separation (PC/H,  $N = 10$ ) and PC without postnatal short-term maternal separation (PC/NH,  $N = 10$ ). All groups were housed in the same animal cages, and all pups were kept together with their mothers. Postnatal short-term maternal separation was performed daily from the 1st to the 14th postnatal day. Separated litters were removed and placed as a group in a plastic cage ( $17.8 \times 28 \times 12.8$  cm) lined with bedding material and placed atop a heating pad maintained at a temperature of  $33 \pm 0.5$  °C (Days 1–5) or  $30 \pm 0.5$  °C (Days 6–14) everyday and returned to their home cages after 15 min (Litvin et al., 2010). Behavioral experiments began when voles were 90 days old.

### Behavioral Testing 1: Open-Field Test

Spontaneous motor activity and anxiety-like behavior were assessed in an open field ( $50 \times 50 \times 50$  cm) illuminated by six 60-W lamps mounted 2 m above the apparatus (Carneiro et al., 2005; Cullen, Burne, Lavidis, & Moritz, 2013; Ermsberger, Azar, & Iwai, 1983). The open field was divided into 25 squares (nine central and 16 peripheral). Adult male offspring were placed individually into the center of the open field for 5 min. The time spent in the central and peripheral zones, total distance covered during the experiment, and the number of crossings between squares were automatically recorded by a digital video system, and anxiety-related indicators analyzed by PC-compatible software (VideoMot 2, TSE Systems, Bad Homburg, Germany) included the frequency and time of entry into the central area, total distance, and number of crossings. The apparatus was cleaned with 70% ethanol after each test, and the interval between experiments was 30 min.

### Behavioral Testing 2: Elevated Plus-Maze Test

After all open-field tests were completed, elevated plus-maze tests were started the next day. The open arm and closed arm of the elevated plus-maze test apparatus were placed 55 cm above the floor with movable pulleys at the bottom. The mandarin voles were moved out from the cage and placed at the connection of the open and closed arms, and their activities in the elevated plus-maze test were recorded by an overhead video camera for 5 min and coded later by a blind observer. The time and frequency of activities in the open and closed arms and the total distance were assessed as indicators of anxiety-related behavior (Lister, 1990; Rodgers & Dalvi, 1997). Voles were returned to their cage when the experiment finished and the apparatus was cleaned with 70% ethanol after each animal.

### Measurement of $E_2$ and T Serum Levels

To avoid the effect of acute-stress-hormone measurement on data, blood samples were collected 2 days after behavior tests, from the anesthetized adult male retro-orbital sinus between 8:00–10:00 a.m. All serum samples were centrifuged from blood at 3,000 rpm for 10 min at room temperature. Serum samples were stored at  $-80$  °C until assays were conducted. Serum T and  $E_2$

concentration assays were measured by using enzyme-linked immunosorbent assay (ELISA; Nieminen et al., 2003). Serum samples were diluted 1:10 to measure T and  $E_2$  levels (He & Tai, 2009). First, the prepared sample and the standard were added to dishes individually and then incubated for 30 min at 37 °C. Second, the dish was washed with solution four times, and a horseradish peroxidase (HRP)-blending agent was added in and incubated for 30 min at 37 °C. Last, after the dish was washed four times, color developing agents A and B were added. After 15 min incubation at 37 °C, the reaction was stopped using a stop solution. The optical density (OD) was determined at 450 nm using a microplate reader (Bio-Tek, Winooski, VT) and taking the blank well as zero. Variation between duplicate values was less than 5%.

### Immunohistochemistry

The brains were collected at the same time as the blood; ER $\alpha$  and AR expression were also examined 2 days after behavioral testing (He, Yu, & Wu, 2013). Voles were deeply anesthetized and perfused with 0.1 molar (M) of phosphate-buffered solution (PBS, pH 7.4) and 4% paraformaldehyde in 0.1 M PBS. The brain was removed within 3 min and placed in 4% paraformaldehyde overnight. Prior to dissection, brains were immersed in 30% sucrose until saturated. Coronal sections (40  $\mu$ m) were cut on a cryostat, and consecutive sections were collected in two vials containing 0.01 M PBS for two different immunohistochemical stainings. The ER $\alpha$  (sc-542; Santa Cruz Biotechnology, Inc., Santa Cruz, CA) and AR antibodies (Ba0004, Neomarkers, Inc., Fremont, CA) are affinity-purified rabbit polyclonal antibodies raised against a peptide and mapped at the C-terminus of mouse origin.

Floating sections were processed using primary antibody and streptavidin and peroxidase methods (Bao & Su, 1991). We incubated the sections for 7 min with 3%  $H_2O_2$  and then washed each of them twice for 10 min with distilled water. We shrank the tissue in 0.01 M PBS. Sections were preincubated for 90 min with normal goat serum (SP-0023, Shanghai Ding Jie Biotechnology, Inc., Shanghai) and incubated at 4 °C overnight with the primary antibody solution (both ER $\alpha$  and AR antibodies, 1:100) diluted by an antibody diluent (0.01 M PBS containing 20% bovine serum albumin and 1.7% Triton X-100: a surfactants). The next day, sections were washed four times for 5 min each with 0.01 M PBS and incubated for 60 min in a 37 °C water bath with biotinylated goat antirabbit antibody (SP-0023), followed by another round of four washes for 5 min each with 0.01 M PBS. After 60 min of incubation with streptavidin/horseradish peroxidase (S-A/HRP) and four washes for 10 min each with 0.01 M PBS, sections were stained with 3,30-diaminobenzidine tetrahydrochloride (DAB) to visualize immunoreactivity. Because ER $\alpha$  and AR were included in nuclei, we counted stained nuclei using an Olympus microscope (Tokyo, Japan).

Slides were randomized and coded for microscopic analysis so that counters were blinded to experimental treatment. The number of cells that showed immunoreactivity was quantified by eye per standard area ( $200 \times 200$   $\mu$ m) using grid sampling. We counted the number of ER $\alpha$ -IRs and AR-IRs in the BNST, mPOA, MeA, PVN, and supraoptic nerve (SON) in 40,000  $\mu$ m<sup>2</sup>. We selected these areas of the brain because they are involved in anxiety (He, Wu, & Yu, 2014; He, Zhang, Shi, & Wang, 2008; Song et al., 2010). Different brain areas were determined according to Nissl-stained brain sections from mandarin voles and a stereotaxic atlas of the rat brain (Bao & Su, 1991).

For each brain nucleus, three representative sections from anterior to posterior and anatomically matched between subjects were chosen and counted to minimize variability. Individual mean values for each animal were obtained by counting positive neurons bilaterally in three sections from each nucleus. Counts were performed separately for each hemisphere, and results were averaged between hemispheres. The left hemisphere was determined from the right hemisphere according to morphological characteristics of the brain surface: After the brain was removed within 3 min, we cut off a small part of the cortex in the left hemisphere and the right hemisphere as a template to discern the left hemisphere from the right hemisphere. Sections were chosen by correspondence to the reference atlas plate instead of the level or intensity of ER $\alpha$ -IR or AR-IR labeling. All immunohistochemistry procedures included negative controls (the primary antibody was not added). A trained experimental rater blinded to experimental treatment counted positive neurons for all subjects. Chosen sections were photographed with a Nikon camera (Tokyo, Japan) attached to a Nikon microscope.

### Statistical Analysis

All data were analyzed using one-sample Kolmogorov-Smirnov tests. These data, including behaviors (time spent, distance traveled, number of crossings, and number of entries), ER $\alpha$ -IRs and AR-IRs in the five brain regions (BNST, MPOA, MeA, PVN and SON), and E<sub>2</sub> and T serum levels were normally distributed, and a two-way factorial analysis of variance (ANOVA) was required to analyze the relationship between mother (PS) and offspring (postnatal short-term maternal separation). If a significant difference was found for all data, post hoc

multiple comparisons were conducted using the Tukey method. The degree of relationship between serum E<sub>2</sub> and T levels and locomotor activity, the distribution of ER $\alpha$ , the time in the center area of the open field, the distribution of AR, and time spent in the open-arm elevated plus-maze test were assessed by calculating correlation coefficients using a two-tailed Pearson product-moment statistic. All data are presented as mean  $\pm$  SE and significance was set at  $p < .05$ . Statistical analyses were conducted using SPSS 10.0 (IBM/SPSS, Chicago, IL).

## Results

### Behavioral Testing 1: Open Field

The two-way ANOVA revealed a significant effect of prenatal handling for time spent in the central area, number of crossings, and total distance. However, postnatal handling did not change significantly for time spent in the central area, number of crossings and total distance. The interaction between both factors was significant for time spent in the central area, number of crossings and total distance. According to post hoc tests, the PS/NH spent less time in the center area, number of crossings and total distance of the open field compared with all other groups ( $p < .05$ ). There were no significant differences between the PC/NH, PC/H, and PS/H groups in spending time in the central area, number of crossings, or total distance in the open open-field test ( $p > .05$ ; Table 1; Figure 1 and 2).

### Behavioral Testing 2: Elevated Plus-Maze Test

Prenatal handling had significant effect on total distance, time spent in the open arm, number of entries to the open arm, and

Table 1  
Main and Interaction F-Statistic Values for the Analyses

| Variable                                       | Two-way ANOVA     |       |                    |      |                                  |       |
|------------------------------------------------|-------------------|-------|--------------------|------|----------------------------------|-------|
|                                                | Prenatal handling |       | Postnatal handling |      | Interaction between both factors |       |
|                                                | F                 | P     | F                  | P    | F                                | P     |
| Time spent in the central area in the OF       | 4.378             | .037  | 2.583              | .081 | 4.579                            | .033  |
| Number of crossings in the OF                  | 4.320             | .038  | 1.889              | .151 | 4.420                            | .036  |
| Total distance in the OF                       | 4.105             | .040  | 2.132              | .095 | 4.632                            | .032  |
| Total distance in the EPM                      | 4.735             | .022  | 2.735              | .073 | 4.645                            | .023  |
| Time spent in the open arms in the EPM         | 4.258             | .038  | 1.978              | .136 | 4.878                            | .015  |
| Number of entries to the open arm in the EPM   | 5.351             | .008  | 1.784              | .152 | 5.251                            | .008  |
| Number of entries to the closed arm in the EPM | 5.155             | .009  | 2.254              | .091 | 4.834                            | .021  |
| E <sub>2</sub> in serum                        | 4.351             | .037  | 2.854              | .069 | 4.843                            | .020  |
| T in serum                                     | 5.758             | .006  | 2.925              | .065 | 5.553                            | .007  |
| ER $\alpha$ -IRs in the MeA                    | 7.767             | .002  | 2.767              | .070 | 8.643                            | <.001 |
| ER $\alpha$ -IRs in the mPOA                   | 9.257             | <.001 | 3.027              | .056 | 9.077                            | <.001 |
| ER $\alpha$ -IRs in the BNST                   | 8.584             | <.001 | 2.784              | .070 | 8.594                            | <.001 |
| ER $\alpha$ -IRs in the PVN                    | 7.952             | <.001 | 2.087              | .121 | 7.967                            | <.001 |
| ER $\alpha$ -IRs in the SON                    | 8.027             | <.001 | 1.781              | .162 | 9.151                            | <.001 |
| AR-IRs in the MeA                              | 6.697             | .003  | 2.871              | .068 | 4.643                            | .023  |
| AR-IRs in the mPOA                             | 8.047             | <.001 | 2.924              | .065 | 17.077                           | <.001 |
| AR-IRs in the BNST                             | 8.013             | <.001 | 2.902              | .066 | 25.594                           | <.001 |

Note. E<sub>2</sub> = estradiol; T = testosterone; ER $\alpha$  = estrogen receptor- $\alpha$ ; AR = androgen receptor; IRs = immunoreactive neurons; OF = open field test; EPM = elevated plus-maze test; PVN = hypothalamic paraventricular nucleus; SON = supraoptic nucleus; BNST = bed nucleus of stria terminalis; mPOA = medial preoptic area; MeA = medial amygdaloid nucleus.




Figure 1. Time spent in the central area and number of crossings in the open-field test. PC/NH: control adult male offspring without postnatal handling; PC/H: control adult male offspring with postnatal handling; PS/NH: prenatal handling without postnatal handling; PS/H: prenatal handling with postnatal handling. The PS/NH spent less time in the center area and made fewer crossings over the open field than all other groups ( $p < .05$ ). There were no significant differences between the PC/NH, PC/H, and PS/H groups in time spent in the central area ( $p > .05$ ) and number of crossings ( $p > .05$ ). All data represent the mean ( $\pm$  SE).

number of entries to the closed arm. Postnatal handling did not significantly change total distance, time spent in the open arm, number of entries to the open arm and number of entries to the closed arm. The interaction between both factors was significant for total distance, time spent in the open arm, number of entries to the open arm, and number of entries to the closed arm. According to post hoc tests, the PS/NH group moved less total distance, spent less time in the open arm, decreased the number of entries to the open arm, but increased number of entries to the closed arm of the elevated plus-maze compared with all other groups ( $p < .05$ ). There were no significant differences between the PC/NH, PC/H, and PS/H groups in moving total distance, time spent in open arms, and number of entries in the open arm and closed arm ( $p > .05$ ; Table 1; Figures 2, 3, and 4).




Figure 2. Total distance in the open field and elevated plus-maze test. The PS/NH moved less total distance of the open field and elevated plus-maze compared with all other groups ( $p < .05$ ). There were no significant differences between the PC/NH, PC/H and PS/H groups in moving total distance of the open field and elevated plus-maze ( $p > .05$ ).




Figure 3. Time spent in the open arm. The PS/NH group spent less time in the open arm of the elevated plus-maze than all other groups. There were no significant differences between the PC/NH, PC/H, and PS/H groups in time spent in the open arm. All data represent the mean ( $\pm$  SE); \*  $p < .05$ .

## E2 and T in Serum

Two-way ANOVA revealed a significant effect of prenatal handling for  $E_2$  and T levels. However, postnatal handling did not change significantly for  $E_2$  levels or T levels. The interaction between both factors was significant for both  $E_2$  and T levels. According to post hoc tests, the serum  $E_2$  level of the PS/NH group was lower than all other groups ( $p < .05$ ), and the serum T level of the PS/NH group was also lower than all other groups ( $p < .05$ ). There were no significant differences between the PC/NH, PC/H, and PS/H groups in  $E_2$  and T serum levels ( $p > .05$ ; Table 1; Figures 5 and 6).




Figure 4. Number of entries into the open and closed arms. The PS/NH group's number of entries into the open arm decreased, but its number of entries into the closed arm of the elevated plus-maze increased compared with all other groups. There were no significant differences between the PC/NH, PC/H, and PS/H groups in the number of entries to the open arm and closed arm. All data represent the mean ( $\pm$  SE); \*  $p < .05$ .




Figure 5. Effects of PS and short-term maternal separation on plasma  $E_2$  levels in adult male offspring. Serum  $E_2$  levels of the PS/NH group were lower than all other groups, and the serum T level of the PS/NH group was also lower compared with all other groups. There were no significant differences between the PC/NH, PC/H, and PS/H groups in  $E_2$  serum levels. All data represent the mean ( $\pm$  SE); \*  $p < .05$ .

### Immunohistochemistry

Prenatal handling had an effect on the number of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN, and SON; postnatal handling did not have such an effect. The interaction between both factors was significant for the number of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN, and SON. The post hoc test showed that the number of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN and SON in the PS/NH group was significantly increased compared with all other groups ( $p < .05$ ). However, there were no significant differences in the number of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN, and SON in the PC/NH compared with the PC/H and PS/H groups in the five brain regions ( $p > .05$ ; Table 1; Figures 7 and 8).




Figure 6. Effects of PS and short-term maternal separation on plasma testosterone levels in adult male offspring. Serum T level of the PS/NH group was lower than all other groups. There were no significant differences between the PC/NH, PC/H, and PS/H groups in T serum levels. All data represent the mean ( $\pm$  SE); \*  $p < .05$ .




Figure 7. Mean ( $\pm$  SE) expression of ER $\alpha$ -IRs after PS and short-term maternal separation. The number of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN, and SON in the PS/NH group was significantly increased compared with all other groups. There were no significant differences between the PC/NH, PC/H, and PS/H groups in the number of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN, and SON. All data represent the mean ( $\pm$  SE); \*  $p < .05$ .

We did not find AR-IRs in the PVN and SON. Two-way ANOVA showed that the prenatal handling affected the number of AR-IRs in the MeA, mPOA, and BNST, whereas postnatal handling did not have the same affect. The interaction between both factors was significant for the number of AR-IRs in the MeA, mPOA, and BNST. The post hoc test showed that the number of AR-IRs in the MeA, mPOA, and BNST in the PS/NH group was significantly reduced compared with all other groups ( $p > .05$ ). However, there were no significant differences in the number of AR-IRs in the MeA, mPOA, or BNST in the PC/NH compared with the PC/H and PS/H groups in the three brain regions ( $p > .05$ ; Table 1; Figures 9 and 10).

### Discussion

#### Prenatal Restraint Stress Increases Anxious Behavior in Adult Male Offspring

In this study, PS males spent less time in the center of the open field and traveled less (Figures 1 and 2). Decreased locomotor activity in these paradigms often reflects increased anxiety rather than some primary effect on locomotion (McKinney et al., 2008). To further assess anxious behavior in PS animals, an elevated plus-maze test, which is a well-validated and conventional test for anxiety-related behavior in animals, was used (Lister, 1987). The result showed that PS animals spent less time engaging in the open arms of the elevated plus-maze test, they made fewer entries into the open arms, and their total distance traveled was lower in the elevated plus-maze test (Figures 2–4). There are data to show that an increase in anxiety is associated with a significant decrease in exploratory behavior in the open arms of the maze, and that the number of entries into the closed arms or the total number of arm entries is a measure of locomotor activity (Shum, Ko, Lee, Kaang,




Figure 8. Position of ER $\alpha$ -IRs in the MeA, mPOA, BNST, PVN, and SON of male voles in the PC/NH, PC/H, PS/NH, and PS/H groups.

& Zhuo, 2005). Thus, all behavior results were indicative of increased anxiety-like behavior: it indicates that, during this stress experience, fetal brain growth is affected and has long-term effects on behaviors. These results are consistent with work on polygynous rat offspring and monogamous guinea pigs (Marrocco et al., 2012; Kapoor & Matthews, 2011) and reveal that prenatal restraint stress is associated with an increased risk of anxiety in adults (McEwen, 2012), regardless of monogamous or polygamous social systems. Previous studies in rats have shown that early PS can cause long-lasting changes in neuroplasticity that result in an increased risk in psychological disorders and stress-related diseases in adult life (Lupien, McEwen, Gunnar, & Heim, 2009). The offspring of pregnant voles receiving daily restraint stress during the last 2 weeks of pregnancy showed decreased serum T levels, and this is consistent with a reduction of serum T levels in rats and guinea pigs in response to high-intensity chronic stimuli (Kapoor & Matthews, 2011).

In normal gestation, male rat fetuses show a surge in T levels on the 18th and 19th gestational days, but such a surge is absent in male fetuses when mothers undergo stress during pregnancy (Ward & Weisz, 1980). Chronic PS increases the level of corticosterone and reduces gonadotropin in serum, inhibiting the activity of the hypothalamic–pituitary–gonadal (HPG) axis and reducing testicular weight (Chen Cárdenas et al., 2013). Stress during gestation can induce long-term effects on the reproductive systems of

male offspring (Arena & Pereira, 2002; Pallarés et al., 2013), thereby affecting their T levels.

We found a negative association between prenatal restraint stress and E<sub>2</sub> levels in serum (see Figure 5), which is inconsistent with our previous work showing a positive association between prenatal treatment and estrogen levels in serum (He, Tai, et al., 2012). This may be because the previous experiment used female offspring and involved ethanol exposure as the stressor. Ethanol is a central nervous system depressant and can cross the placenta and directly affect fetal brain development via several mechanisms. Further, prenatal ethanol stress can increase the aromatization of T, so ethanol increases estrogen levels in serum (He, Tai, et al., 2012; Hilakivi-Clarke, Raygada, & Cho, 1997). However, why prenatal restraint stress decreases serum estrogen levels in adult male offspring is not yet completely understood. It may be that chronic PS inhibits the HPG axis, which reduces plasmatic luteinizing hormone and testicular weight (Chen Cárdenas et al., 2013), thus reducing the secretion of E<sub>2</sub> from testicular cells. PS can also decrease the activity of the steroid aromatase in the male fetal brain (Weinstock, 2005). However, the numbers of ER $\alpha$ -IRs in the BNST, mPOA, MeA, PVN, and SON were enhanced in males exposed to prenatal restraint stress in our study (Figure 7 and 8), and the BNST, MeA, PVN, and SON have been demonstrated to be involved in anxiety (Lee, Macbeth, Pagani, & Young, 2009). ER $\alpha$  is widely expressed throughout most E<sub>2</sub>-sensitive tissues.




Figure 9. Mean ( $\pm$  SE) expression of AR-IRs after PS and short-term maternal separation. The number of AR-IRs in the MeA, mPOA, and BNST in the PS/NH group was significantly reduced compared with all other groups. There were no significant differences between the PC/NH, PC/H, and PS/H groups in the number of AR-IRs in the MeA, mPOA, and BNST. All data represent the mean ( $\pm$  SE); \*  $p < .05$ .

There is differential expression of ER $\alpha$  in some regions, as well as differential brain distribution of changes in ER $\alpha$  that may indicate differential responsiveness to estrogen (Jensen, Jacobson, Walf, & Frye, 2010). Brain regions, including the mPOA, are associated with animal anxiety (Song et al., 2010; He, 2014), and ER $\alpha$  levels are related to anxiety-like behavior (Tsai, Wang, Hong, & Chiu, 2003). ER $\alpha$  activation increases fear and anxiety behavior (Mor-

gan & Pfaff, 2001). In the present experiment, E<sub>2</sub> levels in serum of adult male offspring that underwent PS were lower than males in the PC/NH, PC/H, and PS/H groups, but the number of ER $\alpha$ -IRs in the BNST, mPOA, MeA, PVN, and SON increased (Figures 4, 5, and 6). E<sub>2</sub> levels in serum were lower, and amounts of ER $\alpha$  in brain regions were higher, consistent with other studies (Oliveira et al., 2004). ER $\alpha$  is thought to modulate the activity of corticotropin-releasing factor (CRF). Increased numbers of ER $\alpha$ -IRs in these brain regions raise the possibility that disturbed balance of CRF may affect the activity of the HPA axis and result in anxiety-like behavior (He, Tai, et al., 2012).

We found that serum T levels significantly declined in adult male offspring whose mothers experienced prenatal restraint stress (Figure 6, 9 and 10) and AR-IRs in the MeA, BNST, and mPOA were also fewer in adult male offspring whose mothers experienced prenatal restraint stress. This is consistent with our previous study that decreased T levels in serum and ARs in the central nervous system can result in the development of anxiety-like behavior (He, 2014). The amygdaloid region is a relatively complicated part of the brain and is related to anxiety-like behavior (LeDoux, 2000). It receives information input from the cerebral cortex and subcortical regions and integrates signals from other brain regions including, the BNST, mPOA, PVN, and SON, which ultimately results in expressed anxiousness in animals (Lee et al., 2009). Key brain regions including the BNST, mPOA, PVN, and SON have a high density of AR expression (Williamson & Viau, 2007), and changes in AR-IRs in these areas would likely lead to changes in anxiety. Hence, the significant decline in T levels in serum and the number of AR-IRs in the BNST, mPOA, and MeA increases the risk of anxiety-like behavior in adult male offspring. We speculate that this pattern, i.e., the decrease in T, is likely to cause down regulation of AR in these brain regions and is consis-




Figure 10. Position of AR-IRs in the MeA, mPOA, and BNST of male voles in the PC/NH, PC/H, PS/NH, and PS/H groups.

tent with previous findings that declining T levels in males can decrease or completely eliminate AR-IRs in these brain regions (Arteaga-Silva, Rodríguez-Dorantes, Baig, & Morales-Montor, 2007; He, Zhang, & Guo, 2012). Through ARs, T plays an important role in certain brain regions regulating anxiety-related behavior. For example, anxiety related to T circulation is dependent on ARs (Zuloaga et al., 2008; Zuloaga, Jordan, & Breedlove, 2011) and supports the notion that ARs may play a role in the regulation of anxiety in rodents. However, few AR-IRs were identified in the PVN and SON in our experimental groups. Different distributions of ARs are found across different brain regions and species (Bingham et al., 2006). So far, there is only one report suggesting that prenatal restraint stress may affect AR levels in the mPOA (Ward, Bennett, Ward, Hendricks, & French, 1999), thus, the relationship between the distribution of ARs in other brain regions and prenatal restraint stress is unclear. Prenatal restraint stress can induce long-term effects on the reproductive system of male offspring (Arena & Pereira, 2002; Pallarés et al., 2013). Therefore, prenatal restraint stress may alter AR expression or the HPA axis and affect ARs and T.

### **Postnatal Short-Term Maternal Separation Reverses PS-Induced Anxious Behavior in Adult Male Offspring**

Our open-field test and elevated plus-maze test results show there was no effect of anxiety in the postnatal handling-alone group (Figures 1–4), which is consistent with Lehmann and colleagues' findings that postnatal maternal separation does not affect rat-offspring anxiety (Lehmann et al., 2000). Short periods of separation or handling regimens appear to be beneficial to pups (Lehmann et al., 2000). Although prenatal restraint stress led to reduced motor activity and increased anxiety-like behavior in adult male offspring, and postnatal short-term maternal separation reversed PS-induced anxiety, the result is consistent with findings that increased motor activity and decreased anxiety-like behavior in adult male offspring follow short-term maternal separation (Vallée et al., 1997). In the present test, T levels in serum in the PS/H group were consistent with recent studies demonstrating that postnatal short-term maternal separation induces changes in serum T levels in male rats (Chen Cárdenas et al., 2013). During the critical period of fetal gonadal development, the effect of PS would occur through the reducing activity of aromatase in the hypothalamus (Weinstock, 2005) and 5 alpha-reductase activity in the cerebral cortex and hypothalamus of male fetuses one day after the last session of stress (Ordyan & Pivina, 2005) and D-3- $\beta$ -hydroxysteroid dehydrogenase in the testis (Orth, Weisz, Ward, & Ward, 1983). E2 and T levels in serum in the PS/H group had no significant differences with the PC/NH in the present test that may be occurring by this mechanism which these animals are postnatally stimulated to cause luteinizing hormone and the size of the testis increase and T levels return to normal (Chen Cárdenas et al., 2013). T can be exerted either directly through the AR or indirectly through aromatization to E2 (He, Zhang, et al., 2012). Therefore, E2 and T levels in serum in the PS/H group had no significant differences with the PC/NH. The PS/H voles spent more time, a greater number of crossings, and moved further (i.e., the whole distance) through the open field than the PS/NH group during the test, and moved further than the PS/NH group in the elevated plus-maze test (Figures 1–4). These results are consistent with the

findings that have suggested a positive relationship between plasma testosterone levels and locomotor activity in laboratory-bred male meadow voles (Perrot-Sinal, Innes, Kavaliers, & Ossenkopp, 1998). The presently described relationship between activity and testosterone has been inferred in laboratory-reared animals, mainly on the basis of castration and hormone-replacement manipulations. For instance, injections of testosterone induced significant increases in activity levels in castrated male voles compared with vehicle-injected castrates (Rowsemitt, 1989).

It is known that postnatal short-term maternal separation can cause further changes in an HPA axis already altered by PS, and these changes are likely to influence anxious reactivity (Weinstock, 1997). However, studies on the effects of maternal separation on anxiety have yielded variable results (Kalinichev, Easterling, Plotsky, & Holtzman, 2002) and seem to be related to maternal separation procedures, testing conditions, and stress varieties. For example, adult wild-type offspring fostered by mice with variational biological clocks have been found to exhibit increased anxiety-related behavior (Koizumi, Kurabayashi, Watanabe, & Sanada, 2013). However, in Wistar rats, prolonged maternal separation can lower anxious responses, decrease anxiety, and reverse the effects of adverse conditions on their behavior (Roman, Gustafsson, Berg, & Nylander, 2006). Likewise, pups exposed to chronic unpredictable maternal separation combined with maternal stress develop some resilience to stress when they grow up (Franklin & Mansuy, 2011).

AR has a role in anxiety-like behavior, but the mechanism by which AR activation reduces anxiety is yet to be established. One possibility is that AR stimulation may activate GABAergic drive, known to reduce anxiety (Reynolds, 2008). Here, the distribution of ER $\alpha$ -IRs was also reversed to normal levels in the BNST, mPOA, MeA, SON, and PVN by postnatal short-term maternal separation, which may be related to decreased anxiety-like behavior in adult male offspring.

Postnatal maternal separation causes changes in the mother's behavior to a certain degree, such as increased care of offspring (Levin & Stern, 1975) and increased licking behavior (Barbazanges et al., 1996). This environment-dependent variation in maternal behavior also alters behavioral and HPA responses to stressors in adult offspring (Macri & Würbel, 2006). Mother's behavior may be a result of a number of neuropeptides (e.g., oxytocin and arginine vasopressin), gonadal steroids (e.g., testosterone), and other neurotransmitters being released at parturition and during suckling, promoting the development and maintenance of maternal bonding and maternal care, and further influencing the expression of locomotor activity and exploratory behavioral responses to stress (Curley, Jensen, Mashoodh, & Champagne, 2011).

### **Conclusion**

Despite the limitations in this study, we were able to report the relationship between perinatal stress effects on later anxiety and hormone secretion and ER $\alpha$  and AR in male mandarin voles. In addition, there were some consideration of the possible roles of and changes in ER $\beta$ . As such, we do have another article forthcoming that considers these roles and changes. For the current study, we examined the combined effects of PS and early postnatal maternal separation, which provided insights into the complex relationship between the two. Our data indicated an increase in anxiety-like behaviors in open-field and elevated plus-maze tests

in PS male adults, which was associated with significantly lower estradiol and testosterone levels in serum; an increased number of ER $\alpha$ -IRs in the BNST, mPOA, MeA, PVN, and SON; and a reduced number of AR-IRs in the MeA, mPOA, and BNST. Only early postnatal maternal separation effects were similar to the control group; they did not produce anxiety-like behavior. However, early postnatal maternal separation reversed prenatally induced changes in estradiol and testosterone levels in serum and altered ER $\alpha$ -IRs and AR-IRs in the brain. These data suggest that anxious changes in adults may be governed by early environmental factors and their interaction, because changes in estradiol and testosterone levels in serum and the distribution of ER $\alpha$ -IRs and AR-IRs in the brain result in behavioral changes related to anxiety well into adulthood.

## References

- Arena, A. C., & Pereira, O. C. M. (2002). Neonatal inhalatory anesthetic exposure: Reproductive changes in male rats. *Comparative Biochemistry and Physiology: Part C. Toxicology & Pharmacology*, 133, 633–640. [http://dx.doi.org/10.1016/S1532-0456\(02\)00182-5](http://dx.doi.org/10.1016/S1532-0456(02)00182-5)
- Arpels, J. C. (1996). The female brain hypoestrogenic continuum from the premenstrual syndrome to menopause. A hypothesis and review of supporting data. *The Journal of Reproductive Medicine*, 41, 633–639.
- Arteaga-Silva, M., Rodríguez-Dorantes, M., Baig, S., & Morales-Montor, J. (2007). Effects of castration and hormone replacement on male sexual behavior and pattern of expression in the brain of sex-steroid receptors in BALB/c AnN mice. *Comparative Biochemistry and Physiology: Part A. Molecular & Integrative Physiology*, 147, 607–615. <http://dx.doi.org/10.1016/j.cbpa.2006.11.013>
- Bao, X. M., & Su, S. Y. (1991). *Stereotaxic atlas of the rat brain* (pp. 23–49). Beijing, PRC: People Medical Press.
- Barbazanges, A., Vallée, M., Mayo, W., Day, J., Simon, H., Le Moal, M., & Maccari, S. (1996). Early and later adoptions have different long-term effects on male rat offspring. *The Journal of Neuroscience*, 16, 7783–7790.
- Best, N. R., Rees, M. P., Barlow, D. H., & Cowen, P. J. (1992). Effect of estradiol implant on noradrenergic function and mood in menopausal subjects. *Psychoneuroendocrinology*, 17, 87–93. [http://dx.doi.org/10.1016/0306-4530\(92\)90079-M](http://dx.doi.org/10.1016/0306-4530(92)90079-M)
- Bingham, B., Williamson, M., & Viau, V. (2006). Androgen and estrogen receptor-beta distribution within spinal-projecting and neurosecretory neurons in the paraventricular nucleus of the male rat. *Journal of Comparative Neurology*, 499, 911–923.
- Brake, W. G., Zhang, T. Y., Diorio, J., Meaney, M. J., & Gratton, A. (2004). Influence of early postnatal rearing conditions on mesocortico-limbic dopamine and behavioural responses to psychostimulants and stressors in adult rats. *European Journal of Neuroscience*, 19, 1863–1874. <http://dx.doi.org/10.1111/j.1460-9568.2004.03286.x>
- Carneiro, L. M. V., Diógenes, J. P. L., Vasconcelos, S. M. M., Aragão, G. F., Noronha, E. C., Gomes, P. B., & Viana, G. S. B. (2005). Behavioral and neurochemical effects on rat offspring after prenatal exposure to ethanol. *Neurotoxicology and Teratology*, 27, 585–592. <http://dx.doi.org/10.1016/j.ntt.2005.06.006>
- Carter, C. S., Witt, D. M., Auksi, T., & Casten, L. (1987). Estrogen and the induction of lordosis in female and male prairie voles (*Microtus ochrogaster*). *Hormones and Behavior*, 21, 65–73. [http://dx.doi.org/10.1016/0018-506X\(87\)90031-6](http://dx.doi.org/10.1016/0018-506X(87)90031-6)
- Chen Cárdenas, S. M., Mayer, N., Romanini, M. C., Rolando, A. N., Liaudat, A. C., Brun, N., . . . Rodríguez, N. (2013). Reproductive response in offspring male rats exposed to PS and to early postnatal stimulation. *International Journal of Morphology*, 31, 754–764. <http://dx.doi.org/10.4067/S0717-95022013000200065>
- Cullen, C. L., Burne, T. H. J., Lavidis, N. A., & Moritz, K. M. (2013). Low-dose prenatal ethanol exposure induces anxiety-like behaviour and alters dendritic morphology in the basolateral amygdala of rat offspring. *PLoS ONE*, 8, e54924. <http://dx.doi.org/10.1371/journal.pone.0054924>
- Curley, J. P., Jensen, C. L., Mashoodh, R., & Champagne, F. A. (2011). Social influences on neurobiology and behavior: Epigenetic effects during development. *Psychoneuroendocrinology*, 36, 352–371. <http://dx.doi.org/10.1016/j.psyneuen.2010.06.005>
- Edinger, K. L., & Frye, C. A. (2006). Intrahippocampal administration of an androgen receptor antagonist, flutamide, can increase anxiety-like behavior in intact and DHT-replaced male rats. *Hormones and Behavior*, 50, 216–222. <http://dx.doi.org/10.1016/j.yhbeh.2006.03.003>
- Ellenbroek, B. A., van den Kroonenberg, P. T., & Cools, A. R. (1998). The effects of an early stressful life event on sensorimotor gating in adult rats. *Schizophrenia Research*, 30, 251–260. [http://dx.doi.org/10.1016/S0920-9964\(97\)00149-7](http://dx.doi.org/10.1016/S0920-9964(97)00149-7)
- Ernsberger, P., Azar, S., & Iwai, J. (1983). Open-field behavior in two models of genetic hypertension and the behavioral effects of salt excess. *Behavioral & Neural Biology*, 37, 46–60. [http://dx.doi.org/10.1016/S0163-1047\(83\)91061-0](http://dx.doi.org/10.1016/S0163-1047(83)91061-0)
- Eskelinen, S. I., Vahlberg, T. J., Isoaho, R. E., Kivelä, S. L., & Irjala, K. M. (2007). Associations of sex hormone concentrations with health and life satisfaction in elderly men. *Endocrine Practice*, 13, 743–749. <http://dx.doi.org/10.4158/EP.13.7.743>
- Fedotova, J. (2013). Anxiolytic-like effect of quinpirole in combination with a low dose of 17 $\beta$ -estradiol in ovariectomized rats. *Acta Physiologica Hungarica*, 100, 211–223. <http://dx.doi.org/10.1556/APhysiol.100.2013.2.8>
- Franklin, T. B., & Mansuy, I. M. (2011). The involvement of epigenetic defects in mental retardation. *Neurobiology of Learning and Memory*, 96, 61–67. <http://dx.doi.org/10.1016/j.nlm.2011.04.001>
- Franklin, T. B., Saab, B. J., & Mansuy, I. M. (2012). Neural mechanisms of stress resilience and vulnerability. *Neuron*, 75, 747–761. <http://dx.doi.org/10.1016/j.neuron.2012.08.016>
- Gonzalez, A., Lovic, V., Ward, G. R., Wainwright, P. E., & Fleming, A. S. (2001). Intergenerational effects of complete maternal deprivation and replacement stimulation on maternal behavior and emotionality in female rats. *Developmental Psychobiology*, 38, 11–32. [http://dx.doi.org/10.1002/1098-2302\(2001\)38:1<11::AID-DEV2>3.0.CO;2-B](http://dx.doi.org/10.1002/1098-2302(2001)38:1<11::AID-DEV2>3.0.CO;2-B)
- Granger, D. A., Shirtcliff, E. A., Zahn-Waxler, C., Usher, B., Klimes-Dougan, B., & Hastings, P. (2003). Salivary testosterone diurnal variation and psychopathology in adolescent males and females: Individual differences and developmental effects. *Development and Psychopathology*, 15, 431–449. <http://dx.doi.org/10.1017/S0954579403000233>
- Gregoire, A. J., Kumar, R., Everitt, B., & Studd, J. W. (1996). Transdermal oestrogen for treatment of severe postnatal depression. *The Lancet*, 347, 930–933. [http://dx.doi.org/10.1016/S0140-6736\(96\)91414-2](http://dx.doi.org/10.1016/S0140-6736(96)91414-2)
- Guo, R., Liang, N., Tai, F. D., Wu, R. Y., Chang, G., He, F. Q., & Yuan, Q. W. (2011). Differences in spatial learning and memory for male and female mandarin voles (*Microtus mandarinus*) and BALB/c mice. *Zoological Studies*, 50, 24–30.
- He, F. (2014). The relationship of prenatal ethanol exposure and anxiety-related behaviors and central androgen receptor and vasopressin expression in adult male mandarin voles. *Neuroscience*, 266, 224–234. <http://dx.doi.org/10.1016/j.neuroscience.2014.02.016>
- He, F. Q., & Tai, F. D. (2009). Mating behavior induces changes of expression of Fos protein, plasma testosterone and androgen receptors in the accessory olfactory bulb (AOB) of the male mandarin vole *Microtus mandarinus*. *Current Zoology*, 55, 288–295.
- He, F., Tai, F., Zhang, Y., & Zhang, X. (2012). Effects of castration on aggression and levels of serum sex hormones and their central receptors in mandarin voles (*Microtus mandarinus*). *Journal of Comparative Physiology: A. Neuroethology, Sensory, Neural, and Behavioral Physiology*, 198, 347–362. <http://dx.doi.org/10.1007/s00359-012-0713-6>

- He, F., Wu, R., & Yu, P. (2014). Study of Fos, androgen receptor and testosterone expression in the sub-regions of medial amygdala, bed nucleus of stria terminalis and medial preoptic area in male Mandarin voles in response to chemosensory stimulation. *Behavioural Brain Research*, 258, 65–74. <http://dx.doi.org/10.1016/j.bbr.2013.10.004>
- He, F., Yu, P., & Wu, R. (2013). Relationship between sexual satiety and motivation, brain androgen receptors and testosterone in male mandarin voles. *Behavioural Brain Research*, 250, 257–263. <http://dx.doi.org/10.1016/j.bbr.2013.05.022>
- He, F. Q., Zhang, J., & Guo, X. (2012). Prenatal ethanol exposure increased adult depressive-like behaviors and central ER $\alpha$  and OT expression in female Mandarin voles (*Microtus mandarinus*). *Zoological Studies*, 51, 1–11.
- He, F. Q., Zhang, J. W., Shi, J., & Wang, B. (2008). Changes of estrogen in serum and estrogen receptor  $\beta$  in the relevant brain regions following mating behavior of the male mandarin vole *Microtus mandarinus* [Chinese text]. *Zoological Research*, 29, 529–536.
- Hilakivi-Clarke, L., Raygada, M., & Cho, E. (1997). Serum estradiol levels and ethanol-induced aggression. *Pharmacology, Biochemistry, and Behavior*, 58, 785–791. [http://dx.doi.org/10.1016/S0091-3057\(97\)00035-X](http://dx.doi.org/10.1016/S0091-3057(97)00035-X)
- Howerton, C. L., & Bale, T. L. (2012). Prenatal programming: At the intersection of maternal stress and immune activation. *Hormones and Behavior*, 62, 237–242. <http://dx.doi.org/10.1016/j.yhbeh.2012.03.007>
- Huot, R. L., Plotsky, P. M., Lenox, R. H., & McNamara, R. K. (2002). Neonatal maternal separation reduces hippocampal mossy fiber density in adult Long Evans rats. *Brain Research*, 950, 52–63. [http://dx.doi.org/10.1016/S0006-8993\(02\)02985-2](http://dx.doi.org/10.1016/S0006-8993(02)02985-2)
- Jensen, E. V., Jacobson, H. I., Walf, A. A., & Frye, C. A. (2010). Estrogen action: A historic perspective on the implications of considering alternative approaches. *Physiology & Behavior*, 99, 151–162.
- Kalinichev, M., Easterling, K. W., Plotsky, P. M., & Holtzman, S. G. (2002). Long-lasting changes in stress-induced corticosterone response and anxiety-like behaviors as a consequence of neonatal maternal separation in Long-Evans rats. *Pharmacology, Biochemistry and Behavior*, 73, 131–140. [http://dx.doi.org/10.1016/S0091-3057\(02\)00781-5](http://dx.doi.org/10.1016/S0091-3057(02)00781-5)
- Kapoor, A., & Matthews, S. G. (2011). Testosterone is involved in mediating the effects of PS in male guinea pig offspring. *The Journal of Physiology*, 589, 755–766. <http://dx.doi.org/10.1113/jphysiol.2010.200543>
- Koizumi, H., Kurabayashi, N., Watanabe, Y., & Sanada, K. (2013). Increased anxiety in offspring reared by circadian Clock mutant mice. *PLoS ONE*, 8, e66021. <http://dx.doi.org/10.1371/journal.pone.0066021>
- Krezel, W., Dupont, S., Krust, A., Chambon, P., & Chapman, P. F. (2001). Increased anxiety and synaptic plasticity in estrogen receptor beta-deficient mice. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 12278–12282. <http://dx.doi.org/10.1073/pnas.221451898>
- LeDoux, J. E. (2000). Emotion circuits in the brain. *Annual Review of Neuroscience*, 23, 155–184. <http://dx.doi.org/10.1146/annurev.neuro.23.1.155>
- Lee, H. J., Macbeth, A. H., Pagani, J. H., & Young, W. S., III. (2009). Oxytocin: The great facilitator of life. *Progress in Neurobiology*, 88, 127–151.
- Lehmann, J., Stöhr, T., & Feldon, J. (2000). Long-term effects of PS experiences and postnatal maternal separation on emotionality and attentional processes. *Behavioural Brain Research*, 107, 133–144. [http://dx.doi.org/10.1016/S0166-4328\(99\)00122-9](http://dx.doi.org/10.1016/S0166-4328(99)00122-9)
- Levin, R., & Stern, J. M. (1975). Maternal influences on ontogeny of suckling and feeding rhythms in the rat. *Journal of Comparative and Physiological Psychology*, 89, 711–721. <http://dx.doi.org/10.1037/h0077038>
- Lister, R. G. (1987). The use of a plus-maze to measure anxiety in the mouse. *Psychopharmacology*, 92, 180–185. <http://dx.doi.org/10.1007/BF00177912>
- Lister, R. G. (1990). Ethologically-based animal models of anxiety disorders. *Pharmacology & Therapeutics*, 46, 321–340. [http://dx.doi.org/10.1016/0163-7258\(90\)90021-5](http://dx.doi.org/10.1016/0163-7258(90)90021-5)
- Litvin, Y., Tovote, P., Pentkowski, N. S., Zeyda, T., King, L. B., Vasconcellos, A. J., . . . Blanchard, R. J. (2010). Maternal separation modulates short-term behavioral and physiological indices of the stress response. *Hormones and Behavior*, 58, 241–249. <http://dx.doi.org/10.1016/j.yhbeh.2010.03.010>
- Lovic, V., & Fleming, A. S. (2004). Artificially-reared female rats show reduced prepulse inhibition and deficits in the attentional set shifting task: Reversal of effects with maternal-like licking stimulation. *Behavioural Brain Research*, 148, 209–219. [http://dx.doi.org/10.1016/S0166-4328\(03\)00206-7](http://dx.doi.org/10.1016/S0166-4328(03)00206-7)
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10, 434–445. <http://dx.doi.org/10.1038/nrn2639>
- Macri, S., & Würbel, H. (2006). Developmental plasticity of HPA and fear responses in rats: A critical review of the maternal mediation hypothesis. *Hormones and Behavior*, 50, 667–680. <http://dx.doi.org/10.1016/j.yhbeh.2006.06.015>
- Marrocco, J., Mairesse, J., Ngomba, R. T., Silletti, V., Van Camp, G., Bouwalerh, H., . . . Morley-Fletcher, S. (2012). Anxiety-like behavior of PS rats is associated with a selective reduction of glutamate release in the ventral hippocampus. *The Journal of Neuroscience*, 32, 17143–17154. <http://dx.doi.org/10.1523/JNEUROSCI.1040-12.2012>
- McEwen, B. S. (2012). The ever-changing brain: Cellular and molecular mechanisms for the effects of stressful experiences. *Developmental Neurobiology*, 72, 878–890. <http://dx.doi.org/10.1002/dneu.20968>
- McKinney, B. C., Schneider, J. S., Schafer, G. L., Lowing, J. L., Mohan, S., Zhao, M. X., . . . Murphy, G. G. (2008). Decreased locomotor activity in mice expressing tTA under control of the CaMKII $\alpha$  promoter. *Genes, Brain & Behavior*, 7, 203–213. <http://dx.doi.org/10.1111/j.1601-183X.2007.00339.x>
- Meaney, M. J., Mitchell, J. B., Aitken, D. H., Bhatnagar, S., Bodnoff, S. R., Iny, L. J., & Sarrieau, A. (1991). The effects of neonatal handling on the development of the adrenocortical response to stress: Implications for neuropathology and cognitive deficits in later life. *Psychoneuroendocrinology*, 16, 85–103. [http://dx.doi.org/10.1016/0306-4530\(91\)90072-2](http://dx.doi.org/10.1016/0306-4530(91)90072-2)
- Morgan, M. A., & Pfaff, D. W. (2001). Effects of estrogen on activity and fear-related behaviors in mice. *Hormones and Behavior*, 40, 472–482. <http://dx.doi.org/10.1006/hbeh.2001.1716>
- Morrison, J. H., Brinton, R. D., Schmidt, P. J., & Gore, A. C. (2006). Estrogen, menopause, and the aging brain: How basic neuroscience can inform hormone therapy in women. *The Journal of Neuroscience*, 26, 10332–10348. <http://dx.doi.org/10.1523/JNEUROSCI.3369-06.2006>
- Nestler, E. J., Barrot, M., DiLeone, R. J., Eisch, A. J., Gold, S. J., & Monteggia, L. M. (2002). Neurobiology of depression. *Neuron*, 34, 13–25. [http://dx.doi.org/10.1016/S0896-6273\(02\)00653-0](http://dx.doi.org/10.1016/S0896-6273(02)00653-0)
- Newhouse, P. A., Dumas, J., Wilkins, H., Coderre, E., Sites, C. K., Naylor, M., . . . Young, S. N. (2010). Estrogen treatment impairs cognitive performance after psychosocial stress and monoamine depletion in post-menopausal women. *Menopause: The Journal of the North American Menopause Society*, 17, 860–873. <http://dx.doi.org/10.1097/gme.0b013e3181e15df4>
- Nieminen, P., Mustonen, A. M., Lindström-Seppä, P., Kärkkäinen, V., Mussalo-Rauhamaa, H., & Kukkonen, J. V. K. (2003). Phytosterols affect endocrinology and metabolism of the field vole (*Microtus agrestis*). *Experimental Biology and Medicine*, 228, 188–193.
- Olaso, R., & Habert, R. (2000). Genetic and cellular analysis of male germ cell development. *Journal of Andrology*, 21, 497–511.
- Oliveira, C. A., Mahecha, G. A. B., Carnes, K., Prins, G. S., Saunders, P. T. K., França, L. R., & Hess, R. A. (2004). Differential hormonal regulation of estrogen receptors ER $\alpha$  and ER $\beta$  and androgen receptor expression in rat efferent ductules. *Reproduction: The Journal of the Society for Reproduction and Fertility*, 128, 73–86. <http://dx.doi.org/10.1530/rep.1.00136>

- Ordyan, N. E., & Pivina, S. G. (2005). Effects of PS on the activity of an enzyme involved in neurosteroid synthesis during the "critical period" of sexual differentiation of the brain in male rats. *Neuroscience and Behavioral Physiology*, 35, 931–935. <http://dx.doi.org/10.1007/s11055-005-0148-4>
- Orth, J. M., Weisz, J., Ward, O. B., & Ward, I. L. (1983). Environmental stress alters the developmental pattern of delta 5–3 beta-hydroxysteroid dehydrogenase activity in Leydig cells of fetal rats: A quantitative cytochemical study. *Biology of Reproduction*, 28, 625–631. <http://dx.doi.org/10.1095/biolreprod28.3.625>
- Pallarés, M. E., Adrover, E., Baier, C. J., Bourguignon, N. S., Monteleone, M. C., Brocco, M. A., . . . Antonelli, M. C. (2013). Prenatal maternal restraint stress exposure alters the reproductive hormone profile and testis development of the rat male offspring. *Stress: The International Journal on the Biology of Stress*, 16, 429–440. <http://dx.doi.org/10.3109/10253890.2012.761195>
- Patisaul, H. B., & Bateman, H. L. (2008). Neonatal exposure to endocrine active compounds or an ER $\beta$  agonist increases adult anxiety and aggression in gonadally intact male rats. *Hormones and Behavior*, 53, 580–588. <http://dx.doi.org/10.1016/j.yhbeh.2008.01.008>
- Pelliniemi, L. J., Frojman, K., Paranko, J. (1993). Embryological and prenatal development and function of Sertoli cells. In L. D. Russell & M. D. Griswold (Eds.), *The Sertoli cell* (pp. 88–113). Vienna, IL: Cache River Press.
- Perrot-Sinal, T. S., Innes, D., Kavaliers, M., & Ossenkopp, K. P. (1998). Plasma testosterone levels are related to various aspects of locomotor activity in wild-caught male meadow voles (*Microtus pennsylvanicus*). *Physiology & Behavior*, 64, 31–36. [http://dx.doi.org/10.1016/S0031-9384\(98\)00015-8](http://dx.doi.org/10.1016/S0031-9384(98)00015-8)
- Reynolds, D. S. (2008). The value of genetic and pharmacological approaches to understanding the complexities of GABA(A) receptor subtype functions: The anxiolytic effects of benzodiazepines. *Pharmacology, Biochemistry and Behavior*, 90, 37–42. <http://dx.doi.org/10.1016/j.pbb.2008.03.015>
- Richardson, H. N., Zorrilla, E. P., Mandym, C. D., & Rivier, C. L. (2006). Exposure to repetitive versus varied stress during prenatal development generates two distinct anxiogenic and neuroendocrine profiles in adulthood. *Endocrinology*, 147, 2506–2517. <http://dx.doi.org/10.1210/en.2005-1054>
- Rodgers, R. J., & Dalvi, A. (1997). Anxiety, defence and the elevated plus-maze. *Neuroscience and Biobehavioral Reviews*, 21, 801–810. [http://dx.doi.org/10.1016/S0149-7634\(96\)00058-9](http://dx.doi.org/10.1016/S0149-7634(96)00058-9)
- Roman, E., Gustafsson, L., Berg, M., & Nylander, I. (2006). Behavioral profiles and stress-induced corticosteroid secretion in male Wistar rats subjected to short and prolonged periods of maternal separation. *Hormones and Behavior*, 50, 736–747. <http://dx.doi.org/10.1016/j.yhbeh.2006.06.016>
- Rowsemitt, C. N. (1989). Activity of castrated male voles: Rhythms of responses to testosterone replacement. *Physiology & Behavior*, 45, 7–13. [http://dx.doi.org/10.1016/0031-9384\(89\)90159-5](http://dx.doi.org/10.1016/0031-9384(89)90159-5)
- Sawrey, D. K., & Dewsbury, D. A. (1985). Control of ovulation, vaginal estrus, and behavioral receptivity in voles (*Microtus*). *Neuroscience and Biobehavioral Reviews*, 9, 563–571. [http://dx.doi.org/10.1016/0149-7634\(85\)90003-X](http://dx.doi.org/10.1016/0149-7634(85)90003-X)
- Shum, F. W., Ko, S. W., Lee, Y. S., Kaang, B. K., & Zhuo, M. (2005). Genetic alteration of anxiety and stress-like behavior in mice lacking CaMKIV. *Molecular Pain*, 1, 22. <http://dx.doi.org/10.1186/1744-8069-1-22>
- Sichel, D. A., Cohen, L. S., Robertson, L. M., Rutenber, A., & Rosenbaum, J. F. (1995). Prophylactic estrogen in recurrent postpartum affective disorder. *Biological Psychiatry*, 38, 814–818. [http://dx.doi.org/10.1016/0006-3223\(95\)00063-1](http://dx.doi.org/10.1016/0006-3223(95)00063-1)
- Song, Z., Tai, F., Yu, C., Wu, R., Zhang, X., Broders, H., . . . Guo, R. (2010). Sexual or paternal experiences alter alloparental behavior and the central expression of ER $\alpha$  and OT in male mandarin voles (*Microtus mandarinus*). *Behavioural Brain Research*, 214, 290–300. <http://dx.doi.org/10.1016/j.bbr.2010.05.045>
- Tai, F. D., Wang, T. Z., & Zhao, Y. J. (1999). Study on the breeding and inbreeding avoidance in mandarin vole *Microtus mandarinus* [Chinese text]. *Acta Theriologica Sinica*, 19, 144–149.
- Tai, F. D., Wang, T. Z., & Zhou, Y. J. (2001). Mate choice and related characteristics of Mandarin voles (*Microtus mandarinus*) [Chinese text]. *Acta Zoologica Sinica*, 47, 260–267.
- Tchernitchin, A. N., Tchernitchin, N. N., Mena, M. A., Unda, C., & Soto, J. (1999). Imprinting: Perinatal exposures cause the development of diseases during the adult age. *Acta Biologica Hungarica*, 50, 425–440.
- Tsai, S. J., Wang, Y. C., Hong, C. J., & Chiu, H. J. (2003). Association study of oestrogen receptor alpha gene polymorphism and suicidal behaviours in major depressive disorder. *Psychiatric Genetics*, 13, 19–22. <http://dx.doi.org/10.1097/00041444-200303000-00003>
- Vallée, J. P., Sostman, H. D., MacFall, J. R., Wheeler, T., Hedlund, L. W., Spritzer, C. E., & Coleman, R. E. (1997). MRI quantitative myocardial perfusion with compartmental analysis: A rest and stress study. *Magnetic Resonance in Medicine*, 38, 981–989. <http://dx.doi.org/10.1002/mrm.1910380618>
- Van den Bergh, B. R., Van Calster, B., Smits, T., Van Huffel, S., & Lagae, L. (2008). Antenatal maternal anxiety is related to HPA-axis dysregulation and self-reported depressive symptoms in adolescence: A prospective study on the fetal origins of depressed mood. *Neuropsychopharmacology*, 33, 536–545. <http://dx.doi.org/10.1038/sj.npp.1301450>
- Ward, I. L., Bennett, A. L., Ward, O. B., Hendricks, S. E., & French, J. A. (1999). Androgen threshold to activate copulation differs in male rats prenatally exposed to alcohol, stress, or both factors. *Hormones and Behavior*, 36, 129–140. <http://dx.doi.org/10.1006/hbeh.1999.1534>
- Ward, I. L., & Weisz, J. (1980). Maternal stress alters plasma testosterone in fetal males. *Science*, 207, 328–329. <http://dx.doi.org/10.1126/science.7188648>
- Ward, O. B., Ward, I. L., Denning, J. H., French, J. A., & Hendricks, S. E. (2002). Postparturitional testosterone surge in male offspring of rats stressed and/or fed ethanol during late pregnancy. *Hormones and Behavior*, 41, 229–235. <http://dx.doi.org/10.1006/hbeh.2001.1746>
- Weinstock, M. (1997). Does PS impair coping and regulation of hypothalamic–pituitary–adrenal axis? *Neuroscience and Biobehavioral Reviews*, 21, 1–10. [http://dx.doi.org/10.1016/S0149-7634\(96\)00014-0](http://dx.doi.org/10.1016/S0149-7634(96)00014-0)
- Weinstock, M. (2005). The potential influence of maternal stress hormones on development and mental health of the offspring. *Brain, Behavior, and Immunity*, 19, 296–308. <http://dx.doi.org/10.1016/j.bbi.2004.09.006>
- Williamson, M., & Viau, V. (2007). Androgen receptor expressing neurons that project to the paraventricular nucleus of the hypothalamus in the male rat. *The Journal of Comparative Neurology*, 503, 717–740. <http://dx.doi.org/10.1002/cne.21411>
- Young, K. A., Gobrogge, K. L., Liu, Y., & Wang, Z. (2011). The neurobiology of pair bonding: Insights from a socially monogamous rodent. *Frontiers in Neuroendocrinology*, 32, 53–69. <http://dx.doi.org/10.1016/j.yfrne.2010.07.006>
- Zuloaga, D. G., Jordan, C. L., & Breedlove, S. M. (2011). The organizational role of testicular hormones and the androgen receptor in anxiety-related behaviors and sensorimotor gating in rats. *Endocrinology*, 152, 1572–1581. <http://dx.doi.org/10.1210/en.2010-1016>
- Zuloaga, D. G., Morris, J. A., Jordan, C. L., & Breedlove, S. M. (2008). Mice with the testicular feminization mutation demonstrate a role for androgen receptors in the regulation of anxiety-related behaviors and the hypothalamic–pituitary–adrenal axis. *Hormones and Behavior*, 54, 758–766. <http://dx.doi.org/10.1016/j.yhbeh.2008.08.004>

Received May 3, 2015

Revision received July 26, 2015

Accepted July 30, 2015 ■