

BRIEF COMMUNICATION

Differential Influence of Social Versus Isolate Housing on Vicarious Fear Learning in Adolescent Mice

Jules B. Panksepp and Gareth P. Lahvis
Oregon Health and Science University

Laboratory rodents can adopt the pain or fear of nearby conspecifics. This phenotype conceptually lies within the domain of empathy, a bio-psycho-social process through which individuals come to share each other's emotion. Using a model of cue-conditioned fear, we show here that the expression of vicarious fear varies with respect to whether mice are raised socially or in solitude during adolescence. The impact of the adolescent housing environment was selective: (a) vicarious fear was more influenced than directly acquired fear, (b) "long-term" (24-h postconditioning) vicarious fear memories were stronger than "short-term" (15-min postconditioning) memories in socially reared mice whereas the opposite was true for isolate mice, and (c) females were more fearful than males. Housing differences during adolescence did not alter the general mobility of mice or their vocal response to receiving the unconditioned stimulus. Previous work with this mouse model underscored a genetic influence on vicarious fear learning, and the present study complements these findings by elucidating an interaction between the adolescent social environment and vicarious experience. Collectively, these findings are relevant to developing models of empathy amenable to mechanistic exploitation in the laboratory.

Keywords: empathy, emotion, conditioning, rodent, behavior

Survival and reproduction require that an animal engage in social interactions. Such actions always include mating-related behaviors, but can extend to nurturance of offspring, alloparenting, cooperative defense and hunting, as well as formation and maintenance of hierarchies. Social participants can benefit in such contexts when they are aware of the affective state of others; for instance, if a calm infant becomes agitated, if companions increase vigilance to a threat, or if a peer changes its receptivity toward play solicitations. Empathy, or expressing "an affective response more appropriate to another's situation than to one's own" (Hoffman, 2001), is a mechanism through which emotions can be shared (Preston & de Waal, 2002). Behavioral mimicry is another way by which individuals can learn from others, but does not require emotion, and may be less beneficial than empathy when the social environment is variable or difficult to predict (Nakahashi & Ohtsuki, 2015).

Comparative and evolutionary approaches to understanding empathy (de Waal, 2012; Mogil, 2012; Panksepp & Lahvis, 2011; Panksepp & Panksepp, 2013) indicate that it manifests as a compilation of subprocesses (Preston & de Waal, 2002). In this regard studies have demonstrated that laboratory mice express such "endophenotypes" (Chen, Panksepp, & Lahvis, 2009; Jeon et al.,

2010; Langford et al., 2006; Sanders, Mayford, & Jeste, 2013; Gonzalez-Liencre, Juckel, Tas, Friebe, & Brüne, 2014). For example, observer C57BL/6J mice exhibit vicarious behavioral (Chen et al., 2009; Jeon et al., 2010), physiological (Chen et al., 2009), and neural responses (Jeon et al., 2010) to other's expressions of fear. These studies and others using laboratory rats (Atsak et al., 2011; Ben-Ami Bartal, Rodgers, Bernardez Sarria, Decety, & Mason, 2014; Jones, Riha, Gore, & Monfils, 2014) additionally support the notion that social relationships, sexual characteristics, stress, and prior experience are potent modulators of empathic responsiveness (Christov-Moore, Simpson, Coudé, Grigaityte, Iacoboni, & Ferrari, 2014; Freidin, Carballo, & Bentosela, 2015; Martin et al., 2015).

Social exclusion can also influence empathy (Twenge, Baumeister, DeWall, Ciarocco, & Bartels, 2007), particularly during adolescent development (Eisenberg, 1982). In rodents, manipulation of the social housing environment is used as an experimental procedure to restrict or augment social interaction during adolescence, and can profoundly affect maturation of the brain and behavior (Adams & Rosenkranz, 2015; Gan, Bowline, Lourenco, & Pickel, 2014; Liu et al., 2012; Makinodan, Rosen, Ito, & Corfas, 2012; Yang, Perry, Weber, Katz, & Crawley, 2011). In the present study, we hypothesized that social housing during mouse adolescence would have a supportive effect on vicarious fear relative to isolate housing. We compared males and females because sexual identity can influence empathic responding. Moreover, responses were assessed 15-min and 24-h postconditioning to distinguish between "short-term" and "long-term" memories, respectively, the latter of which may be less sensitive to the acute experience associated with conditioning.

This article was published Online First February 15, 2016.

Jules B. Panksepp and Gareth P. Lahvis, Department of Behavioral Neuroscience, Oregon Health and Science University.

Correspondence concerning this article should be addressed to Jules B. Panksepp, Department of Behavioral Neuroscience, Oregon Health and Science University, 3181 SW Sam Jackson Park Road, L470, Portland, OR 97239-3011. E-mail: panksepp@ohsu.edu

Method

Mice from the BALB/cJ (“BALB”) and C57BL/6J (“B6”) mice were bred within our own colony. At weaning B6 mice were pooled from 2–4 litters (Figure 1A), and then randomly selected

for housing in either a social group of 2 males and 2 females (see Panksepp et al., 2007 for rationale) or in complete social isolation (Figure 1B). These B6 mice then became observers of target mice undergoing fear conditioning (i.e., vicarious fear) or were directly




Figure 1. Animal husbandry timeline, behavioral procedure, and the effects of differential social housing on the expression of vicarious and directly acquired fear. (A) Litters of B6 juveniles were raised in their natal litter without the sire. (B) Juveniles were then weaned into either mixed-sex groups or social isolation where they were housed throughout the experiment. (C) B6 observers experienced unfamiliar F1 targets undergo cued-fear conditioning whereas B6 targets were directly conditioned. These experimental groups were compared with control groups that did not directly experience the US. (D) B6 observers and targets were then tested for the expression of a fear response as a function of differential social housing at 15-min and 24-h postconditioning. (E) Isolate and socially housed mice expressed equivalent vicarious fear responses at 15-min, but this phenotype decreased and increased, respectively, at 24-h (see Results). (F) Isolate mice exhibited longer freezing responses than socially housed mice at 15-min postconditioning, but this difference was reversed at 24-h. *Ns* for (E) and (F) are provided within each respective bar. Control mice from the ISO and SOC groups were pooled in (E) and (F) as there were no differences between them (statistics not shown). Note that the ordinates for the vicarious and direct groups are on different scales. Data are presented as the mean $\pm$ SE. Asterisks represent significant orthogonal contrasts. * $p < .05$. ** $p < .01$. *** $p < .001$. # $p < .05$ for comparisons between the 15-min and 24-h time points.

conditioned (see Figure 1C and below). Mice remained in their respective housing conditions throughout conditioning and testing. To control for familiarity with strain-specific communication modalities, such as vocalizations or volatile odorants, target mice during vicarious fear conditioning were age matched male-female pairs of unfamiliar mice derived from reciprocal BALB $\times$ B6 (F1) crosses (see Chen et al., 2009 for rationale).

On Day 1, B6 observers and targets were allowed 300-s to freely explore the fear-conditioning compartment of the experimental apparatus (www.cleversysinc.com/products/hardware). To familiarize observers with a distressful stimulus, they were exposed to a single, unconditioned stimulus (US; 3-s, 1 mA scrambled shock) halfway through this habituation period (see Chen et al., 2009 for rationale; also see Sanders et al., 2013 for empirical support). Note that directly conditioned B6 targets did not receive US pre-exposure, but were habituated to the conditioning compartment. Exposure of observers to the US during the habituation period did not change subsequent baseline freezing when re-exposed to the conditioning chamber for testing (see Results), indicating that this experience did not engender contextual fear (for an example of vicarious fear using contextual conditioning see Jeon et al., 2010). The inefficacy of US pre-exposure to produce contextual freezing during testing is consistent with previous findings (Chen et al., 2009), and may be partially accounted for by the fact that the “observation” chambers and conditioning compartment shared a common feature (i.e., stainless steel dowels; see below), which may lead to partial extinction of contextual fear that results from US pre-exposure. Additionally, observers received a single US during habituation and were handled by the experimenter during the intervening cued-conditioning protocols (see below), which are aspects of this procedure that deviate from more traditional contextual-conditioning paradigms and may explain why US pre-exposure in observers did not increase baseline freezing during testing.

Fifteen-min after the habituation period B6 observers and targets were subjected to vicarious and direct cued-fear conditioning, respectively (Figure 1C). During vicarious conditioning, two observers (a male and a female) were isolated individually into observation chambers adjacent to the conditioning compartment and separated from each other by opaque Plexiglas. The floor and one wall of the observation chambers were composed of inactive stainless steel dowels, identical to the active dowels of the conditioning compartment on the floor and on the wall facing the observation chambers. Conditioning and testing was conducted under infrared LED arrays during the middle 6-h of the dark phase when laboratory mice are generally awake and active, so observers could hear and smell F1 targets, but could not touch or see them. Dim red lighting in the colony and testing room during experimental procedures was ≈ 15 lux.

After 120-s in the conditioning compartment pairs of F1 targets used for vicarious fear conditioning, or an individual B6 target for direct fear conditioning, were exposed to 10 presentations of a 30-s auditory stimulus (1 kHz pure tone, 85 dB) that coterminated with the US (3-s, 1 mA scrambled shock). US-conditional stimulus (CS) pairings were separated by 90-s intervals. The conditioning protocol was repeated on Day 2. Although this conditioning protocol is somewhat extreme for the direct conditioning groups, it is used for vicarious conditioning to elicit robust activation of social communication modalities between targets and observers (e.g., see

Jeon et al., 2010). Mice in control groups were treated exactly as described above except that F1 targets and B6 targets were not exposed to the US (see Figure 1C). Vocalizations at frequencies below 150 kHz were recorded during all conditioning sessions on Day 2 with the UltraSoundGate 416H recording system and CM16 condenser microphones (Avisoft Bioacoustics).

Testing entailed placing an individual B6 observer or target into the conditioning compartment 15-min (Day 2) and 24-h (Day 3) after the second conditioning session (Figure 1D). These testing time points were selected because of their standard use as respective short-term and long-term assays of classical conditioned memory. After 120-s in the conditioning compartment each mouse was exposed to five presentations of the CS (each spaced by a 90-s interval). All components of the experimental apparatus were thoroughly cleaned with 70% ethanol and water, and dried between each phase of the experiment. Administration of all conditioning stimuli was controlled automatically (FreezeScan, Cleversys Inc.). Infrared video cameras recorded mouse behaviors during the test sessions. Freezing was defined as the complete absence of movement other than respiratory movements and was assessed with computer-assisted software (ButtonBox v.5.0, Behavioral Research Solutions) for the duration of each CS presentation (i.e., cued fear) and for the 60-s leading up to the first CS (i.e., pre-cue, baseline fear). Each measurement was repeated by the first author—blinded to the experimental condition—and all data presentation and statistical outcomes were based on the average of these two measurements. “Intrarater” reliability was high for both cued (Pearson’s correlation, $R = 0.99$, $df = 1,289$) and pre-CS measurements ($R = 0.99$, $df = 257$). A subset of the cued trials (59% of the data) were also evaluated for interrater reliability by comparing the measurements of the first author to those of a blinded laboratory technician ($R = 0.98$, $df = 765$). Distress vocalizations (DVs) during US administration (see Chen et al., 2009) were tallied using the “interacting labeling” function in the SASLab Pro software package (Avisoft Bioacoustics). DVs were identified by their characteristic resonant energies above the CS and above background noise. For each conditioning session on Day 2, DVs were quantified two times in a blinded fashion by the first author ($R = 0.91$, $df = 57$) and averaged across the session. Data presentation and statistical outcomes were based on the average of these two measurements. The “automatic parameters” function was also used to extract the fundamental frequency (pitch), amplitude and duration of each DV.

Results

Vicariously conditioned mice expressed higher levels of CS-induced freezing relative to control mice when they were tested at the 15-min short-term time point (Figure 1E; $F(2, 327) = 10.5$, $p < .0001$), but there was no difference between isolate and socially housed mice (orthogonal contrast, $F(1, 324) < 0.1$, $p = .96$). A main effect of conditioning was also found 24-h after the last vicarious conditioning trial, $F(2, 327) = 30.2$, $p < .0001$. Compared with 15-min postconditioning, CS-induced freezing was more sensitive to housing conditions of mice at this time point: Socially housed mice expressed higher levels of freezing than isolate mice (orthogonal contrast, $F(1, 324) = 30.3$, $p < .0001$), a difference attributable to a decrease in isolated mice (two-tailed paired t tests; $t = 12.6$, $p = .01$, $df = 218$) and an

Table 1
Freezing Responses of Observers and Target Mice 24-h
Postconditioning as a Function of Housing Condition and Sex

Housing-type	Conditioning-type	Sex	% Freezing
Isolate	Direct	M	67.9 ± 2.23
Social	Direct	M	63.9 ± 3.40
Isolate	Direct	F	53.0 ± 2.88
Social	Direct	F	70.0 ± 2.49*
Isolate	Vicarious	M	7.8 ± 1.74
Social	Vicarious	M	11.5 ± 1.33
Isolate	Vicarious	F	6.2 ± 1.16
Social	Vicarious	F	18.8 ± 2.26*

Note. Freezing responses of directly and vicariously conditioned B6 mice are displayed relative to their housing context and sex. Data are presented as the mean ± SE. Asterisks represent significant orthogonal contrasts comparing isolate versus social housing contexts-by-sex for each type of conditioning.

* $p < .001$.

increase in socially housed mice ($t = 12.81$, $p = .02$, $df = 238$) relative to the 15-min time point. Isolated mice expressed marginally higher levels of CS-induced freezing relative to their controls at the 24-h time point (orthogonal contrast, $F(1, 324) = 4.1$, $p = .04$).

For mice receiving vicarious conditioning CS-induced freezing was not sensitive to their sexual identity at either time point nor was there a housing-by-sex interaction at 15-min postconditioning (statistics not shown). However, a housing-by-sex interaction was found at the 24-h time point, $F(2, 327) = 5.4$, $p < .01$. Males (Table 1; orthogonal contrast, $F(1, 324) = 3.1$, $p = .08$) were much less influenced by the housing conditions than females, $F(1, 324) = 36.3$, $p < .0001$ when tested 24-h after the last vicarious conditioning experience.

Mice conditioned by direct exposure to the US-CS pairing exhibited robust CS-induced freezing responses compared with control mice, $F(2, 297) = 225.3$, $p < .0001$. Fifteen-min after conditioning isolate mice were more likely to freeze than socially housed mice (Figure 1F; orthogonal contrast, $F(1, 294) = 9.1$, $p < .01$). Twenty-four hours after direct fear conditioning mice were also substantially more likely to freeze than their respective controls, $F(2, 307) = 532.7$, $p < .0001$. However, while isolate mice were more likely to freeze than socially housed mice when tested 15-min after conditioning, they exhibited less CS-induced freezing than socially housed mice when tested 24-h postconditioning (Figure 1F; orthogonal contrast, $F(1, 304) = 7.0$, $p < .01$).

Similar to mice conditioned vicariously, the sexual identity of mice did not affect freezing 15-min or 24-h after direct conditioning nor was there a housing-by-sex interaction detected when mice were tested 15-min postconditioning (statistics not shown). A housing-by-sex interaction was present 24-h after direct fear conditioning, $F(2, 307) = 11.4$, $p < .0001$, with socially housed females expressing higher levels of freezing than their isolated counterparts (Table 1; $F(1, 304) = 25.6$, $p < .0001$). By contrast, isolated males and socially housed males exhibited similar levels of CS-induced freezing, $F(1, 304) = 1.6$, $p = .21$ after direct fear conditioning.

Because socially housed females from both the vicarious and direct groups expressed higher levels of freezing than isolated females 24-h postconditioning, we used the average freezing time

of isolate females as a baseline to assess the magnitude to which social housing affected the vicarious and direct fear phenotypes. There was a larger difference between isolate and socially housed females that underwent vicarious conditioning (mean change from isolation ± SE, +213.5 ± 37.65%) compared with females with directly acquired fear (+32.0 ± 4.69%; $F(1, 108) = 19.1$, $p < .0001$).

Social influences on CS-induced freezing were not attributable to factors unrelated to the US-CS pairing during vicarious or direct conditioning. For instance, differences in pre-CS freezing were not detected for vicariously conditioned mice relative to controls at the short-term or long-term testing time points (Figure 2A; both $ps > 0.10$). Although pre-CS freezing was higher at both time points for mice that were directly conditioned relative to controls (Figure 2B; both $ps < 0.0001$), there were no differences in pre-CS freezing between directly conditioned isolate and socially housed mice at either postconditioning time point (both $ps > 0.25$).

The number of DVs emitted during direct conditioning did not differ as a function of social housing (Table 2; $F(1, 33) = 1.8$, $p = .19$) nor did DV pitch, amplitude or duration differ between isolate and socially housed mice (all $ps > 0.55$). Effects of sex or housing-by-sex interactions were not detected for any of the DV variables (statistics not shown). DV emission, pitch, amplitude, and duration were also similar between target F1 mice used for vicarious conditioning of isolate and socially housed observer mice (all $ps > 0.54$).

Discussion

Empathy is a complex social ability mediated by interactions between several subprocesses, but it is fundamentally governed by emotional substrates. Advances in understanding the biological underpinnings of empathy have come from careful laboratory studies of rodents (for a review see Panksepp & Lahvis, 2011; Panksepp & Panksepp, 2013) that complement those in other




Figure 2. Precue freezing after placement into the conditioning chamber following vicarious or direct fear conditioning. (A) Baseline freezing upon placement into the conditioning chamber was minimal after vicarious conditioning and did not differ between groups at 15-min or 24-h postconditioning. (B) Increased baseline freezing responses after direct fear conditioning were detected, and were similar in isolate and socially housed mice at both time points. Note that the ordinates for the vicarious and direct groups are on different scales. Data are presented as the mean ± SE. Asterisks represent significant orthogonal contrasts. *** $p < .001$.

Table 2
Distress Vocalizations of Fear Conditioned Mice

Housing-type	Conditioning-type	DVs
Isolate	Direct	9.5 ± .65
Social	Direct	8.6 ± .29
Isolate	Vicarious	12.0 ± .91*
Social	Vicarious	12.5 ± .70*

Note. The number of DVs emitted by directly conditioned B6 mice and F1 target-pairs from vicarious conditioning are displayed. Note that DVs in the vicarious condition represent the acoustic emissions of 2 mice versus 1 during direct conditioning. Data are presented as the mean ± *SE*. Asterisks represent significant Tukey-Kramer post hoc tests comparing direct versus vicarious conditioning for each housing context.

* *p* < .05.

species (Decety, 2011). In this respect, we described the (short-term) vicarious fear phenotype in adolescent mice over 6 years ago at our former laboratory site, with a different batch of B6 mice, and using a slightly modified conditioning procedure (Chen et al., 2009), which suggests this behavioral response is relatively stable despite potential sources of uncontrolled variability.

Our current findings provide further evidence that this mouse vicarious fear phenotype models some basic features of empathy. In humans, strong attachment to peers during adolescence can be influential for empathic responding (Laible, Carlo, & Raffaelli, 2000) and female empathy moderates social relationships with peers (Laible, Carlo, & Roesch, 2004). Consistent with these studies, we found adolescent mice restricted from social interactions (and social relationships with cage mates) express blunted vicarious fear 24-h postconditioning and females were more affected than male observers. The finding that vicarious fear in female mice was more sensitive to the adolescent housing environment than males is consistent with the idea that sex differences in empathy likely have deep phylogenetic and ontogenetic underpinnings (Christov-Moore, Simpson, Coudé, Grigaityte, Iacoboni, & Ferrari, 2014).

Had social isolation induced changes in stress reactivity to conspecific exposure, we would have expected to find group differences in freezing shortly after (i.e., 15-min) vicarious conditioning, but we did not. By contrast isolate mice expressed higher levels of freezing at the short-term testing time point when they were targets indicating that social isolation increases fear acquisition via direct conditioning. Furthermore, isolation did not alter the general mobility of mice, as there were no housing-based differences in baseline freezing before CS administration during testing. The emission of DVs (i.e., “audible squeaks”) to the US during conditioning, which communicates fear to observers (Chen et al., 2009) and is an index of sensitivity to US administration in rodents (e.g., Ji, Fu, Adwanikar, & Neugebauer, 2013)—was similarly unchanged by social versus isolate housing. Collectively these findings indicate that social contact can enhance the learning abilities of adolescent female mice particularly when the task is focused on the emotional state of others.

The vicarious fear phenotype was increased and decreased in the social versus isolate housing conditions, respectively, only 24-h after conditioning, which raises questions about the events that occur between conditioning and the long-term testing time point. After conditioning, socially reared mice may communicate with

each other via changes in ultrasonic vocalizations, odorants and/or behavioral activity in the home cage, and this could foster the expression of vicarious fear. By contrast, restriction from such supportive influences during social isolation after conditioning could result in a reduction of vicarious fear. Growing up in a social group versus solitude may also influence processes associated with memory consolidation, an interpretation that we favor because of the selectivity of the adolescent housing environment on affecting vicarious fear 24-h (but not 15-min) postconditioning. Such hypotheses regarding the time-sensitivity of social versus isolate housing on expression of the vicarious fear phenotype can be further explored experimentally. Nevertheless, adolescent mice developing in a group presumably have more social competence than those raised in isolation (e.g., Liu et al., 2012; Makinodan et al., 2012). If this is the case, isolated mice would likely be more influenced in the vicarious conditioning paradigm, which involves social communication, than by direct conditioning, which does not include a social component, as was found here.

It is especially noteworthy that “observational fear” between adolescent rats, but not directly acquired fear, is similarly affected by social versus isolate housing during adolescence (Yusufshaq & Rosenkranz, 2013), which suggests that across species this period of development is important for the maturation of empathy. Whether adolescence constitutes a “critical period” for emergence of the capacity for vicarious fear remains an open issue. Moreover, the distinct pattern of increased/decreased fear across housing contexts, conditioning procedures and testing times that we observed beckons an additional point to consider: These differential and interactive influences on CS-induced freezing suggest socially acquired fear may have biological mechanisms (genomic, synaptic, or neuro-anatomical) different from those involved in directly acquired fear, which may have important implications for understanding the neurobiology of empathy in animals.

References

- Adams, T., & Rosenkranz, J. A. (2015). Social isolation during postweaning development causes hypoactivity of neurons in the medial nucleus of the male rat amygdala. [Advance online publication]. *Neuropsychopharmacology*. <http://dx.doi.org/10.1038/npp.2015.364>
- Atsak, P., Orre, M., Bakker, P., Cerliani, L., Roozendaal, B., Gazzola, V., . . . Keysers, C. (2011). Experience modulates vicarious freezing in rats: A model for empathy. *Public Library of Science One*, 6, e21855. <http://dx.doi.org/10.1371/journal.pone.0021855>
- Ben-Ami Bartal, I., Rodgers, D. A., Bernardez Sarria, M. S., Decety, J., & Mason, P. (2014). Pro-social behavior in rats is modulated by social experience. *eLife*, 3, e01385.
- Chen, Q., Panksepp, J. B., & Lahvis, G. P. (2009). Empathy is moderated by genetic background in mice. *Public Library of Science One*, 4, e4387. <http://dx.doi.org/10.1371/journal.pone.0004387>
- Christov-Moore, L., Simpson, E. A., Coudé, G., Grigaityte, K., Iacoboni, M., & Ferrari, P. F. (2014). Empathy: Gender effects in brain and behavior. *Neuroscience and Biobehavioral Reviews*, 46, 604–627. <http://dx.doi.org/10.1016/j.neubiorev.2014.09.001>
- Decety, J. (2011). The neuroevolution of empathy. *Annals of the New York Academy of Sciences*, 1231, 35–45. <http://dx.doi.org/10.1111/j.1749-6632.2011.06027.x>
- de Waal, F. B. (2012). The antiquity of empathy. *Science*, 336, 874–876. <http://dx.doi.org/10.1126/science.1220999>
- Eisenberg, N. (1982). *The development of prosocial behavior*. New York, NY: Academic Press.

- Freidin, E., Carballo, F., & Bentosela, M. (2015). Direct reciprocity in animals: The roles of bonding and affective processes. [Advance online publication]. *International Journal of Psychology*. <http://dx.doi.org/10.1002/ijop.12215>
- Gan, J. O., Bowline, E., Lourenco, F. S., & Pickel, V. M. (2014). Adolescent social isolation enhances the plasmalemmal density of NMDA NR1 subunits in dendritic spines of principal neurons in the basolateral amygdala of adult mice. *Neuroscience*, 258, 174–183. <http://dx.doi.org/10.1016/j.neuroscience.2013.11.003>
- Gonzalez-Lieners, C., Juckel, G., Tas, C., Friebe, A., & Brüne, M. (2014). Emotional contagion in mice: The role of familiarity. *Behavioural Brain Research*, 263, 16–21. <http://dx.doi.org/10.1016/j.bbr.2014.01.020>
- Hoffman, M. (2001). *Empathy and moral development*. Cambridge, United Kingdom: Cambridge University Press.
- Jeon, D., Kim, S., Chetana, M., Jo, D., Ruley, H. E., Lin, S. Y., . . . Shin, H. S. (2010). Observational fear learning involves affective pain system and Cav1.2 Ca²⁺ channels in ACC. *Nature Neuroscience*, 13, 482–488. <http://dx.doi.org/10.1038/nn.2504>
- Ji, G., Fu, Y., Adwanikar, H., & Neugebauer, V. (2013). Non-pain-related CRF1 activation in the amygdala facilitates synaptic transmission and pain responses. *Molecular Pain*, 9, 2. <http://dx.doi.org/10.1186/1744-8069-9-2>
- Jones, C. E., Riha, P. D., Gore, A. C., & Monfils, M.-H. (2014). Social transmission of Pavlovian fear: Fear-conditioning by-proxy in related female rats. *Animal Cognition*, 17, 827–834. <http://dx.doi.org/10.1007/s10071-013-0711-2>
- Laible, D. J., Carlo, G., & Raffaelli, M. (2000). The differential relations of parent and peer attachment to adolescent adjustment. *Journal of Youth and Adolescence*, 29, 45–59. <http://dx.doi.org/10.1023/A:1005169004882>
- Laible, D. J., Carlo, G., & Roesch, S. C. (2004). Pathways to self-esteem in late adolescence: The role of parent and peer attachment, empathy, and social behaviours. *Journal of Adolescence*, 27, 703–716. <http://dx.doi.org/10.1016/j.adolescence.2004.05.005>
- Langford, D. J., Cragger, S. E., Shehzad, Z., Smith, S. B., Sotocinal, S. G., Levenstadt, J. S., . . . Mogil, J. S. (2006). Social modulation of pain as evidence for empathy in mice. *Science*, 312, 1967–1970. <http://dx.doi.org/10.1126/science.1128322>
- Liu, J., Dietz, K., DeLoyht, J. M., Pedre, X., Kelkar, D., Kaur, J., . . . Casaccia, P. (2012). Impaired adult myelination in the prefrontal cortex of socially isolated mice. *Nature Neuroscience*, 15, 1621–1623. <http://dx.doi.org/10.1038/nn.3263>
- Makinodan, M., Rosen, K. M., Ito, S., & Corfas, G. (2012). A critical period for social experience-dependent oligodendrocyte maturation and myelination. *Science*, 337, 1357–1360. <http://dx.doi.org/10.1126/science.1220845>
- Martin, L. J., Hathaway, G., Isbester, K., Mirali, S., Acland, E. L., Niederstrasser, N., . . . Mogil, J. S. (2015). Reducing social stress elicits emotional contagion of pain in mouse and human strangers. *Current Biology*, 25, 326–332. <http://dx.doi.org/10.1016/j.cub.2014.11.028>
- Mogil, J. S. (2012). The surprising empathic abilities of rodents. *Trends in Cognitive Sciences*, 16, 143–144. <http://dx.doi.org/10.1016/j.tics.2011.12.012>
- Nakahashi, W., & Ohtsuki, H. (2015). When is emotional contagion adaptive? *Journal of Theoretical Biology*, 380, 480–488. <http://dx.doi.org/10.1016/j.jtbi.2015.06.014>
- Panksepp, J. B., Jochman, K. A., Kim, J. U., Koy, J. J., Wilson, E. D., Chen, Q., . . . Lahvis, G. P. (2007). Affiliative behavior, ultrasonic communication and social reward are influenced by genetic variation in adolescent mice. *Public Library of Science One*, 2, e351. <http://dx.doi.org/10.1371/journal.pone.0000351>
- Panksepp, J. B., & Lahvis, G. P. (2011). Rodent empathy and affective neuroscience. *Neuroscience and Biobehavioral Reviews*, 35, 1864–1875. <http://dx.doi.org/10.1016/j.neubiorev.2011.05.013>
- Panksepp, J., & Panksepp, J. B. (2013). Toward a cross-species understanding of empathy. *Trends in Neurosciences*, 36, 489–496. <http://dx.doi.org/10.1016/j.tins.2013.04.009>
- Preston, S. D., & de Waal, F. B. (2002). Empathy: Its ultimate and proximate bases. *Behavioral and Brain Sciences*, 25, 1–20.
- Sanders, J., Mayford, M., & Jeste, D. (2013). Empathic fear responses in mice are triggered by recognition of a shared experience. *Public Library of Science One*, 8, e74609. <http://dx.doi.org/10.1371/journal.pone.0074609>
- Twenge, J. M., Baumeister, R. F., DeWall, C. N., Ciarocco, N. J., & Bartels, J. M. (2007). Social exclusion decreases prosocial behavior. *Journal of Personality and Social Psychology*, 92, 56–66. <http://dx.doi.org/10.1037/0022-3514.92.1.56>
- Yang, M., Perry, K., Weber, M. D., Katz, A. M., & Crawley, J. N. (2011). Social peers rescue autism-relevant sociability deficits in adolescent mice. *Autism Research*, 4, 17–27. <http://dx.doi.org/10.1002/aur.163>
- Yusufishaq, S., & Rosenkranz, J. A. (2013). Post-weaning social isolation impairs observational fear conditioning. *Behavioural Brain Research*, 242, 142–149. <http://dx.doi.org/10.1016/j.bbr.2012.12.050>

Received October 19, 2015

Revision received January 3, 2016

Accepted January 5, 2016 ■